

Twitter Thread by Walter M Chesnut



Walter M Chesnut

@Parsifaler



1) THERE IS A VERY REAL, CREDIBLE AND SIGNIFICANT DANGER THAT mRNA SPIKE PROTEIN THERAPIES WILL INDUCE PRION OR PRION-LIKE DISEASE

A paper from 2016 shows that prion and prion-like proteins can also spread and propagate from cell to cell through macropinocytosis. It is also

2) noted that the Spike Protein of SARS-CoV-2 can enter cells through macropinocytosis, without the presence of the entire virion. Given the recent Tweet to me by @endocdoc, where he recently saw a patient develop Creutzfeld-Jakob after COVID-19, the extant paper showing CJD

3) CJD after COVID-19 infection and the recent death of the head of Astra-Zeneca research from CJD should be concerning to all. Especially if he gave himself the AZ therapy during Stage 1 or Stage 2 trials. This would give a possible timeframe for onset.

Again, all Spike Protein

4) mRNA and DNA should be paused until long term animal studies can be conducted.

<https://t.co/hBBqN1hyDO>

<https://t.co/M1pANLHXHJ>

<https://t.co/LsQ6GJGTNY>

<https://t.co/oPdLu4z5vZ>

5)

<https://t.co/PWOgyPoG5d>

<https://t.co/dE8tsNgvaj>

