Twitter Thread by Walter M Chesnut



Walter M Chesnut
@Parsifaler

1) THERE IS A VERY REAL, CREDIBLE AND SIGNIFICANT DANGER THAT mRNA SPIKE PROTEIN THERAPIES WILL INDUCE PRION OR PRION-LIKE DISEASE

A paper from 2016 shows that prion and prion-like proteins can also spread and propagate from cell to cell through macropinocytosis. It is also

2) noted that the Spike Protein of SARS-CoV-2 can enter cells through macropinocytosis, without the presence of the entire virion. Given the recent Tweet to me by @endocdoc, where he recently saw a patient develop Creutzfeld-Jakob after COVID-19, the extant paper showing CJD

3) CJD after COVID-19 infection and the recent death of the head of Astra-Zeneca research from CJD should be concerning to all. Especially if he gave himself the AZ therapy during Stage 1 or Stage 2 trials. This would give a possible timeframe for onset.

Again, all Spike Protein

4) mRNA and DNA should be paused until long term animal studies can be conducted.

https://t.co/hBBqN1hyDO

https://t.co/M1pANLHXHJ


https://t.co/LsQ6GJGTNY

https://t.co/oPdLu4z5vZ

5)

https://t.co/PWOgyPoG5d

https://t.co/dE8tsNgvaj


A
20 µm
mock
MHV
B
% cells with internalized nanoparticles
50
40
30
20
10
0
**
**
mock
MHV
mock
SARS-CoV
DBT
DBT ACE2
C
% cells with internalized nanoparticles
80
60
40
20
0
**
***
UV
HI
MHV
0-2
2-4
4-6
6-8
Nanoparticle addition, h p.i.
D
20 µm
0:01
18:15
34:00
48:45